

Detecting interactions between dark matter and photons at high energy e^+e^- colliders

Zhao-Huan YU (余钊煥)

Institute of High Energy Physics, CAS

with Qi-Shu YAN and Peng-Fei YIN

[arXiv:1307.5740](https://arxiv.org/abs/1307.5740)



Dalian, August 23, 2013

DM-photon interaction

In general, dark matter (DM) are not luminous



DM particles (χ) should not have electric charge
and not directly couple to photons


DM-photon interaction

In general, dark matter (DM) are not luminous



DM particles (χ) should not have electric charge
and not directly couple to photons

However, DM particles may couple to photons via loop diagrams




DM-photon interaction

In general, dark matter (DM) are not luminous



DM particles (χ) should not have electric charge
and not directly couple to photons

However, DM particles may couple to photons via loop diagrams



For **nonrelativistic** DM particles, the photons produced in $\chi\chi \rightarrow \gamma\gamma$ would be **mono-energetic**





A γ -ray line at energy $\sim m_\chi$
("smoking gun" for DM particles)

A γ -ray line from the Galactic center region?

Using the 3.7-year Fermi-LAT γ -ray data, several analyses showed that there might be evidence of **a monochromatic γ -ray line at energy ~ 130 GeV**, originating from the Galactic center region (about $3 - 4\sigma$).

It may be due to DM annihilation with $\langle \sigma_{\text{ann}} v \rangle \sim 10^{-27} \text{ cm}^3 \text{ s}^{-1}$.





Su & Finkbeiner, 1206.1616

Weniger, 1204.2797

Recently, the Fermi-LAT Collaboration has released its official spectral line search in the energy range $5 - 300$ GeV using 3.7 years of data.


They **did not find any globally significant lines** and set 95% CL upper limits for DM annihilation cross sections.

Their most significant fit occurred at $E_\gamma = 133$ GeV and had **a local significance of 3.3σ** , which translates to a global significance of 1.6σ .



Fermi-LAT Collaboration, 1305.5597

DM-photon interaction at e^+e^- colliders



The coupling between DM particles and photons that induce the annihilation process $\chi\chi \rightarrow \gamma\gamma$ can also lead to the process $e^+e^- \rightarrow \chi\chi\gamma$. Therefore, the possible γ -ray line signal observed by Fermi-LAT may be tested at future TeV-scale e^+e^- colliders.

DM particles escape from the detector



Signature: a **monophoton** associating with missing energy ($\gamma + \cancel{E}$)

Effective operator approach

If DM particles couple to photons via exchanging some mediators which are **sufficiently heavy**, the DM-photon coupling can be approximately described by **effective contact operators**.

For Dirac fermionic DM, consider $\mathcal{O}_F = \frac{1}{\Lambda^3} \bar{\chi} i\gamma_5 \chi F_{\mu\nu} \tilde{F}^{\mu\nu}$:

$$\langle \sigma_{\text{ann}} v \rangle_{\chi\bar{\chi} \rightarrow 2\gamma} \simeq \frac{4m_\chi^4}{\pi\Lambda^6}, \quad \sigma(e^+e^- \rightarrow \chi\bar{\chi}\gamma) \sim \frac{s^2}{\Lambda^6}$$


Fermi γ -ray line signal $\iff m_\chi \simeq 130 \text{ GeV}, \Lambda \sim 1 \text{ TeV}$

For complex scalar DM, consider $\mathcal{O}_S = \frac{1}{\Lambda^2} \chi^* \chi F_{\mu\nu} F^{\mu\nu}$:

$$\langle \sigma_{\text{ann}} v \rangle_{\chi\chi^* \rightarrow 2\gamma} \simeq \frac{2m_\chi^2}{\pi\Lambda^4}, \quad \sigma(e^+e^- \rightarrow \chi\chi^*\gamma) \sim \frac{s}{\Lambda^4}$$

Fermi γ -ray line signal $\iff m_\chi \simeq 130 \text{ GeV}, \Lambda \sim 3 \text{ TeV}$

In the $\gamma + \cancel{E}$ searching channel, the main background is $e^+e^- \rightarrow \nu\bar{\nu}\gamma$:






Minor backgrounds: $e^+e^- \rightarrow e^+e^-\gamma$, $e^+e^- \rightarrow \tau^+\tau^-\gamma$, ...

Simulation: FeynRules → MadGraph 5 → PGS 4

$$\text{ILD-like ECAL energy resolution: } \frac{\Delta E}{E} = \frac{16.6\%}{\sqrt{E/\text{GeV}}} \oplus 1.1\%$$




Future e^+e^- colliders: $\sqrt{s} = 250 \text{ GeV}$ ("Higgs factory"),
 $\sqrt{s} = 500 \text{ GeV}$ (typical ILC), $\sqrt{s} = 1 \text{ TeV}$ (upgraded ILC & initial CLIC),
 $\sqrt{s} = 3 \text{ TeV}$ (ultimate CLIC)






Cut 1 (pre-selection):

Require a photon with $E_\gamma > 10$ GeV
and $10^\circ < \theta_\gamma < 170^\circ$
Veto any other particle

Benchmark point: $\Lambda = 200$ GeV, $m_\chi = 100(50)$ GeV for fermionic (scalar) DM



Benchmark point: $\Lambda = 200$ GeV, $m_\chi = 100(50)$ GeV for fermionic (scalar) DM



Cut 1 (pre-selection):




Require a photon with $E_\gamma > 10$ GeV and $10^\circ < \theta_\gamma < 170^\circ$

Veto any other particle

Cut 2: Veto $50 \text{ GeV} < m_{\text{miss}} < 130 \text{ GeV}$

Cut 3: Require $30^\circ < \theta_\gamma < 150^\circ$

Benchmark point: $\Lambda = 200$ GeV, $m_\chi = 100(50)$ GeV for fermionic (scalar) DM



Cut 1 (pre-selection):

Require a photon with $E_\gamma > 10$ GeV and $10^\circ < \theta_\gamma < 170^\circ$

Veto any other particle

Cut 2: Veto $50 \text{ GeV} < m_{\text{miss}} < 130 \text{ GeV}$

Cut 3: Require $30^\circ < \theta_\gamma < 150^\circ$

Cut 4: Require $p_T^\gamma > \sqrt{s}/10$

Benchmark point: $\Lambda = 200$ GeV, $m_\chi = 100(50)$ GeV for fermionic (scalar) DM

Cross sections and signal significances after each cut

	$\nu\bar{\nu}\gamma$	$e^+e^-\gamma$	Fermionic DM	Scalar DM		
	σ (fb)	σ (fb)	σ (fb)	S/\sqrt{B}	σ (fb)	S/\sqrt{B}
Cut 1	2415.2	173.0	646.8	12.7	321.4	6.3
Cut 2	2102.5	168.6	646.8	13.6	308.2	6.5
Cut 3	1161.1	16.8	538.0	15.7	255.9	7.5
Cut 4	254.5	1.9	520.7	32.5	253.9	15.8



Benchmark point: $\Lambda = 200$ GeV, $m_\chi = 100(50)$ GeV for fermionic (scalar) DM

Most of the signal events remain

$e^+e^- \rightarrow \nu\bar{\nu}\gamma$ background: reduced by almost **an order of magnitude**

$e^+e^- \rightarrow e^+e^-\gamma$ background: only **one percent** survives

$$(\sqrt{s} = 500 \text{ GeV}, 1 \text{ fb}^{-1})$$






Solid lines: 100 fb^{-1} ; dot-dashed lines: 1000 fb^{-1} ($S/\sqrt{B} = 3$)

ILC luminosity: $240 - 570 \text{ fb}^{-1}/\text{year}$ [ILC TDR, Vol. 1, 1306.6327]

Beam polarization



For a process at an e^+e^- collider with **polarized beams**,

$$\sigma(P_{e^-}, P_{e^+}) = \frac{1}{4} [(1 + P_{e^-})(1 + P_{e^+})\sigma_{RR} + (1 - P_{e^-})(1 - P_{e^+})\sigma_{LL} + (1 + P_{e^-})(1 - P_{e^+})\sigma_{RL} + (1 - P_{e^-})(1 + P_{e^+})\sigma_{LR}]$$



$\blacktriangle (P_{e^-}, P_{e^+}) = (0.8, -0.3)$ can be achieved at the ILC

[ILC technical design report, Vol. 1, 1306.6327]



$$(S/\sqrt{B} = 3)$$

Using the **polarized beams** is roughly equivalent to **increasing** the integrated luminosity by **an order of magnitude**.

For fermionic DM (scalar DM), a data set of 2000 fb^{-1} (1000 fb^{-1}) would be just sufficient to test the Fermi γ -ray line signal at an e^+e^- collider with $\sqrt{s} = 1 \text{ TeV}$ (3 TeV).

S-matrix unitarity

For quantum scattering theories,

S -matrix unitarity ($S^\dagger S = 1$) \Leftrightarrow conservation of probability

A process violate the unitarity in a non-renormalizable effective theory



The theory is **invalid** for this process



A **UV-complete theory** may be needed for a full description

The effective operator treatment for DM searches at colliders should be carefully checked by verifying the S -matrix unitarity.

Unitarity conditions

The $2 \rightarrow 2$ amplitude $\mathcal{M}(\cos \theta)$ can be expanded as **partial waves**:

$$\mathcal{M}(\cos \theta) = 16\pi \sum_j (2j+1) a_j P_j(\cos \theta), \quad a_j = \frac{1}{32\pi} \int_{-1}^1 d \cos \theta P_j(\cos \theta) \mathcal{M}(\cos \theta)$$

Unitarity condition for $2 \rightarrow 2$ **elastic scattering**:

$$|\operatorname{Re} a_j^{\text{el}}| \leq \frac{1}{2}, \quad \forall j$$

Unitarity condition for $2 \rightarrow 2$ **inelastic scattering**:

$$|a_j^{\text{inel}}| \leq \frac{1}{2\sqrt{\beta_f}}, \quad \forall j$$

(β_f is the velocity of either of the final particles)

$$S^\dagger S = 1, S = 1 + iT \Rightarrow -i(T - T^\dagger) = T^\dagger T$$



$$-i(\mathcal{M}_{\alpha \rightarrow \beta} - \mathcal{M}_{\beta \rightarrow \alpha}^*) = \sum_{\gamma} \int d\Pi_{\gamma} \mathcal{M}_{\beta \rightarrow \gamma}^* \mathcal{M}_{\alpha \rightarrow \gamma} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma})$$



$$\begin{aligned} 2 \operatorname{Im} \mathcal{M}_{\text{el}}(\cos \theta_{\alpha \beta}) &= \int d\Pi_{\gamma_{\text{el}}} \mathcal{M}_{\beta \rightarrow \gamma_{\text{el}}}^* \mathcal{M}_{\alpha \rightarrow \gamma_{\text{el}}} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma_{\text{el}}}) \\ &\quad + \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma_n}) + \text{other inelastic terms} \\ &\geq \frac{1}{32\pi^2} \int d\Omega_{k_1} \mathcal{M}_{\text{el}}^*(\cos \theta_{\beta \gamma}) \mathcal{M}_{\text{el}}(\cos \theta_{\alpha \gamma}) + \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma_n}) \end{aligned}$$



$$\operatorname{Im} a_j^{\text{el}} \geq |a_j^{\text{el}}|^2 + |b_j^{\text{inel}}|^2,$$


$$|b_j^{\text{inel}}|^2 \equiv \frac{1}{64\pi} \int d\cos \theta_{\alpha \beta} P_j(\cos \theta_{\alpha \beta}) \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma_n})$$



Unitarity condition for any $2 \rightarrow n$ **inelastic scattering**:



$$|b_j^{\text{inel}}| \leq \frac{1}{2}, \quad \forall j$$

Unitarity bounds: $2 \rightarrow 2$ vs $2 \rightarrow 3$





Given the same \sqrt{s} , unitarity bounds for $2 \rightarrow 2$ scattering are **much more stringent** than those for $2 \rightarrow 3$ scattering.

However, here the relevant bounds are those for $2 \rightarrow 3$ scattering.



All the experimental reaches we obtained lie far beyond the unitarity violation regions.





All the experimental reaches we obtained lie far beyond the unitarity violation regions.

From the viewpoint of S -matrix unitarity, our effective operator treatment do not exceed its valid range.

Conclusions and discussions

- ① In this work, we explore the sensitivity to **the effective operators of DM and photons** at TeV-scale e^+e^- colliders.
- ② With a 100 fb^{-1} dataset, the potential Fermi γ -ray line signal for **the fermionic DM can be tested** at a 3 TeV collider, though **the scalar DM searching would be challenging**.
- ③ Using the **polarized beams** is roughly equivalent to **collecting 10 times of data**.
- ④ In order to check the validity of the effective operator approach, we derive **a general unitarity condition for $2 \rightarrow n$ processes**. **The experimental reaches we obtained are valid** since they lie far beyond the unitarity violation regions.

- ⑤ The unitarity condition for $2 \rightarrow n$ scattering can be also applied to other interesting processes, e.g., the $WW\gamma$ and $WZ\gamma$ production induced by **anomalous quartic gauge couplings**.



[CMS PAS SMP-13-009]

Motivations
○○○○

Sensitivity
○○○○○

Beam polarization
○○

Unitarity bounds
○○○○○

Conclusions
○○●

Backups
○○○○○○○

Thanks for your attentions!

Motivations
○○○○

Sensitivity
○○○○○

Beam polarization
○○

Unitarity bounds
○○○○○

Conclusions
○○○

Backups
●○○○○○○

Backup slides

Note that our unitarity condition $|b_j^{\text{inel}}| \leq \frac{1}{2}$ is derived without any approximation.

Through an approximate method, a unitarity bound on the $2 \rightarrow n$ inelastic cross section $\sigma_{\text{inel}}(2 \rightarrow n)$ can be derived to be

$$\sigma_{\text{inel}}(2 \rightarrow n) \leq \frac{4\pi}{s}.$$


[Dicus & H. -J. He, hep-ph/0409131]

We have compared the results given by these two formulas and find that **their differences are rather small** for the processes considered here.

Unitarity condition in terms of amplitudes:

$$-i(\mathcal{M}_{\alpha \rightarrow \beta} - \mathcal{M}_{\beta \rightarrow \alpha}^*) = \sum_{\gamma} \int d\Pi_{\gamma} \mathcal{M}_{\beta \rightarrow \gamma}^* \mathcal{M}_{\alpha \rightarrow \gamma} (2\pi)^4 \delta^{(4)}(p_{\alpha} - p_{\gamma})$$

For the elastic process $1 + 2 \rightarrow 1 + 2$, consider the transitions of state:



Since $\mathcal{M}_{\alpha \rightarrow \beta} = \mathcal{M}_{\beta \rightarrow \alpha}^* = \mathcal{M}_{\text{el}}(\cos \theta_{\alpha\beta})$, the unitarity condition becomes

$$\begin{aligned} & 2 \operatorname{Im} \mathcal{M}_{\text{el}}(\cos \theta_{\alpha\beta}) \\ = & \int d\Pi_{\gamma_{\text{el}}} \mathcal{M}_{\beta \rightarrow \gamma_{\text{el}}}^* \mathcal{M}_{\alpha \rightarrow \gamma_{\text{el}}} (2\pi)^4 \delta^{(4)}(p_\alpha - p_{\gamma_{\text{el}}}) + \text{inelastic terms} \\ \geq & \frac{\beta_1}{32\pi^2} \int d\Omega_{k_1} \mathcal{M}_{\text{el}}^*(\cos \theta_{\beta\gamma}) \mathcal{M}_{\text{el}}(\cos \theta_{\alpha\gamma}), \end{aligned}$$

where $\beta_1 \equiv \sqrt{1 - 4m_1^2/s}$ and $d\Omega_{k_1} = d\phi_{k_1} d\cos \theta_{\alpha\gamma}$.

In terms of **partial waves**:

$$\begin{aligned} \operatorname{Im} a_j^{\text{el}} & \geq \frac{\beta_1}{8\pi} \sum_{k,l} (2k+1)(2l+1) a_k^{\text{el}*} a_l^{\text{el}} \int d\cos \theta_{\alpha\beta} d\Omega_{k_1} \\ & \quad \times P_j(\cos \theta_{\alpha\beta}) P_k(\cos \theta_{\beta\gamma}) P_l(\cos \theta_{\alpha\gamma}) \end{aligned}$$

The **addition theorem** for Legendre polynomials:

$$P_k(\cos \theta_{\beta\gamma}) = P_k(\cos \theta_{\alpha\beta})P_k(\cos \theta_{\alpha\gamma}) + 2 \sum_{m=1}^l \frac{(l-m)!}{(l+m)!} P_k^m(\cos \theta_{\alpha\beta})P_k^m(\cos \theta_{\alpha\gamma}) \cos m\phi_{k_1}$$

Carrying out all the integrations, we have


$$\text{Im } a_j^{\text{el}} \geq \beta_1 |a_j^{\text{el}}|^2,$$

which is equivalent to

$$(\text{Re } a_j^{\text{el}})^2 + \left(\text{Im } a_j^{\text{el}} - \frac{1}{2\beta_1} \right)^2 \leq \frac{1}{(2\beta_1)^2}.$$


For the scattering of massless particles, $\beta_1 = 1$, and it implies

$$|\text{Re } a_j^{\text{el}}| \leq \frac{1}{2}, \quad \forall j.$$




For $2 \rightarrow n$ inelastic scattering, consider the transitions of state:


$$\alpha(p_1, p_2) \rightarrow \beta(q_1, q_2)$$



$$\alpha(p_1, p_2) \rightarrow \gamma_n(k_3, \dots, k_{n+2})$$



$$\beta(q_1, q_2) \rightarrow \gamma_n(k_3, \dots, k_{n+2})$$



The unitarity condition becomes

$$\begin{aligned} 2 \operatorname{Im} \mathcal{M}_{\text{el}}(\cos \theta_{\alpha\beta}) &= \int d\Pi_{\gamma_{\text{el}}} \mathcal{M}_{\beta \rightarrow \gamma_{\text{el}}}^* \mathcal{M}_{\alpha \rightarrow \gamma_{\text{el}}} (2\pi)^4 \delta^{(4)}(p_\alpha - p_{\gamma_{\text{el}}}) \\ &\quad + \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_\alpha - p_{\gamma_n}) + \text{other inelastic terms} \\ &\geq \frac{\beta_1}{32\pi^2} \int d\Omega_{k_1} \mathcal{M}_{\text{el}}^*(\cos \theta_{\beta\gamma}) \mathcal{M}_{\text{el}}(\cos \theta_{\alpha\gamma}) + \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_\alpha - p_{\gamma_n}). \end{aligned}$$

Introducing a new quantity

$$|b_j^{\text{inel}}|^2 \equiv \frac{1}{64\pi} \int d\cos \theta_{\alpha\beta} P_j(\cos \theta_{\alpha\beta}) \int d\Pi_{\gamma_n} \mathcal{M}_{\beta \rightarrow \gamma_n}^* \mathcal{M}_{\alpha \rightarrow \gamma_n} (2\pi)^4 \delta^{(4)}(p_\alpha - p_{\gamma_n}),$$

we have $\operatorname{Im} a_j^{\text{el}} \geq \beta_1 |a_j^{\text{el}}|^2 + |b_j^{\text{inel}}|^2$. Thus

$$|b_j^{\text{inel}}|^2 \leq \frac{1}{4\beta_1} - \beta_1 \left[(\operatorname{Re} a_j^{\text{el}})^2 + \left(\operatorname{Im} a_j^{\text{el}} - \frac{1}{2\beta_1} \right)^2 \right] \leq \frac{1}{4\beta_1}.$$

For massless incoming particles,

$$|b_j^{\text{inel}}| \leq \frac{1}{2}, \quad \forall j.$$